


Senegal


Total population (2016): 15 589 485 ▶ Population aged 15 years and older (15+): 56% ▶ Population in urban areas: 44% ▶ Income group (World Bank): Low income

ALCOHOL CONSUMPTION: LEVELS AND PATTERNS

Recorded alcohol per capita (15+) consumption, 1961–2016



Recorded alcohol per capita (15+) consumption (in litres of pure alcohol) by type of alcoholic beverage, 2016 or latest year available



Alcohol per capita (15+) consumption (in litres of pure alcohol)

	2010*		2016*	
Recorded	0.3		0.2	
Unrecorded	0.3		0.5	
Total**	0.6		0.7	
Total males / females	1.0	0.2	1.3	0.2
WHO African Region	6.3		6.3	

* Three-year averages of recorded and unrecorded for 2009–2011 and 2015–2017; **adjusted for tourist consumption.

Total alcohol per capita (15+) consumption, drinkers only (in litres of pure alcohol), 2016

	Litres
Males (15+)	4.4
Females (15+)	1.6
Both sexes (15+)	3.5

Prevalence of heavy episodic drinking* (%), 2016

	Population (15+ years)	Drinkers only (15+ years)	Population (15–19 years)	Drinkers only (15–19 years)
Males	12.6	41.6	8.0	44.1
Females	1.7	13.3	1.0	14.4
Both sexes	6.9	32.9	4.5	36.2

* Consumed at least 60 grams or more of pure alcohol on at least one occasion in the past 30 days.

Abstainers (%), 2016

	Males	Females	Both sexes
Lifetime abstainers (15+)	64.4	83.4	74.3
Former drinkers* (15+)	5.2	4.1	4.6
Abstainers (15+), past 12 months	69.6	87.5	78.9

* Persons who used to drink alcoholic beverages but have not done so in the past 12 months.

HEALTH CONSEQUENCES: MORTALITY AND MORBIDITY

Age-standardized death rates (ASDR) and alcohol-attributable fractions (AAF), 2016

	ASDR*		AAF (%)		AAD** (Number)
Liver cirrhosis, males / females	46.8	27.4	14.7	10.5	253
Road traffic injuries, males / females	74.8	29.0	13.3	9.2	546
Cancer, males / females	104.0	98.8	1.6	0.6	60

* Per 100 000 population (15+); **alcohol-attributable deaths, both sexes.

Years of life lost (YLL) score*, 2016

LEAST < 1 **2** 3 4 5 > MOST

* Based on alcohol-attributable years of life lost.

Prevalence of alcohol use disorders and alcohol dependence (%), 2016*

	Alcohol use disorders**	Alcohol dependence
Males	1.4	1.4
Females	0.0	0.0
Both sexes	0.7	0.7
WHO African Region	3.7	1.3

* 12-month prevalence estimates (15+); **including alcohol dependence and harmful use of alcohol.

POLICIES AND INTERVENTIONS

Written national policy (adopted/revised) / National action plan	No / —
Excise tax on beer / wine / spirits	Yes / Yes / Yes
National legal minimum age for off-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
National legal minimum age for on-premise sales of alcoholic beverages (beer / wine / spirits)	18 / 18 / 18
Restrictions on on-/off-premise sales of alcoholic beverages (any): Hours, days / places, density Specific events / intoxicated persons / petrol stations	Yes, No / Yes, Yes Yes / Yes / No

National maximum legal blood alcohol concentration (BAC) when driving a vehicle (general / young / professional), in %	No / No / No
Legally binding regulations on alcohol advertising / product placement (any)	Yes / Yes
Legally binding regulations on alcohol sponsorship / sales promotion (any)	Yes / Yes
Legally required health warning labels on alcohol advertisements / containers (any)	Yes / Yes
National government support for community action (any)	Yes
National monitoring system(s) (any)	No